

ISI B.Stat. (Hons.) & B.Math. (Hons.) 2013 Subjective Solution Paper

$$\begin{aligned}
 1. \quad S &= \log_a bc + \log_b ac + \log_c ba \\
 &= \log_a b + \log_a c + \log_b c + \log_b a + \log_c a + \log_c b \\
 &= \frac{\log b}{\log a} + \frac{\log c}{\log a} + \frac{\log c}{\log b} + \frac{\log a}{\log b} + \frac{\log a}{\log c} + \frac{\log b}{\log c} \\
 &= \left(\frac{\log b}{\log a} + \frac{\log a}{\log b} \right) + \left(\frac{\log c}{\log a} + \frac{\log a}{\log c} \right) + \left(\frac{\log c}{\log b} + \frac{\log b}{\log c} \right)
 \end{aligned}$$

Now applying AM \geq GM inequalities we get,

$$\geq 2 + 2 + 2 = 6.$$

$$2. \quad f'(x) = \frac{2\sin x - 1}{(x + 2\cos x)^2} = 0 \text{ gives } x = \frac{\pi}{6}.$$

$$f(0) = \frac{1}{2}; \quad f\left(\frac{\pi}{6}\right) = \frac{1}{\frac{\pi}{6} + \sqrt{3}} \approx 0.44$$

The function is decreasing in the interval $\left[0, \frac{\pi}{6}\right)$

$$\text{Now, } f\left(\pi - \frac{\pi}{6}\right) = \frac{1}{\frac{5\pi}{6} + \sqrt{3}} \approx 1.13$$

For x in $\left[\frac{\pi}{6}, \frac{5\pi}{6}\right]$, $f(x)$ is positive & then again it becomes decreasing function.

So, the range of the function is $\left(0, \frac{6}{5\pi - 6\sqrt{3}}\right]$.


3. Given that

$$\begin{aligned}
 & -y^2 \leq f(x+y) - f(x-y) - y \leq y^2 \\
 \Rightarrow & -y^2 + y \leq f(x+y) - f(x-y) \leq y^2 + y \\
 \Rightarrow & \frac{-y^2 + y}{(x+y) - (x-y)} \leq \frac{f(x+y) - f(x-y)}{(x+y) - (x-y)} \leq \frac{y^2 + y}{(x+y) - (x-y)} \\
 \Rightarrow & \lim_{2y \rightarrow 0} \left(-\frac{y}{2} + \frac{1}{2}\right) \leq \lim_{2y \rightarrow 0} \frac{f(x+y) - f(x-y)}{(x+y) - (x-y)} \leq \lim_{2y \rightarrow 0} \left(\frac{y}{2} + \frac{1}{2}\right) \\
 \Rightarrow & \frac{1}{2} \leq \lim_{2y \rightarrow 0} \frac{f(x+y) - f(x-y)}{(x+y) - (x-y)} \leq \frac{1}{2}
 \end{aligned}$$

So, $f(x)$ is a linear function $y = ax + c$, where $a = \frac{1}{2}$ & c is a constant.

4. Suppose there is no such player. Say X be the player with highest number of names in his list. Let A be the set of players in list of X whom X has directly defeated and B be the set of other players of X's list. Since there is no player with all others names, say X does not have Y's name. This implies Y has beaten X and hence in Y's list, there is X and also each member of set A. Now the members of set B are in X's list because each is beaten by someone of set A. Now if a member of set A beats Y, that would imply that Y is in X's list, which is not so. Hence Y has beaten each member of set A, which further implies he has the names of set B also in his list. Thus Y is a player with more names in his list than that of X, a contradiction.

5.



Let $\angle AOB = \angle BOC = m$

$$\cos m = \frac{OA^2 + OB^2 - AB^2}{2 \times OA \times OB} = \frac{r^2 + r^2 - \frac{r^2}{4}}{2r^2} = \frac{7}{8}$$

$$\text{Then } \cos 2m = 2\cos^2 m - 1 = \frac{34}{64}$$

Using Cosine Rule in $\triangle AOC$, we get $AC^2 = OA^2 + OC^2 - 2 \times OA \times OC \cos 2m$

$$\text{So, } AC^2 = \frac{2r^2 \times 30}{64}$$

$$\text{Now, } CD^2 = AD^2 - AC^2 = \frac{2r^2 \times 98}{64}$$

$$\text{So, } CD = \frac{7r}{4}$$

$$\Leftrightarrow \frac{CD}{r} = \frac{7}{4}$$

6.

Assume $P(x) - Q(x) = F(x)$. We will show that $F(x)$ has repeated roots of $x = 1$. To prove this we will show that the derivatives of $F(x)$ also has roots $x = 1$.

$$(P(x))^3 - (Q(x))^3 = P(x^3) - Q(x^3) .$$

$$\Rightarrow (P - Q)^3 + 3PQ(P - Q) = P(x^3) - Q(x^3)$$

$$\Rightarrow (F(x))^3 + 3PQF(x) = F(x^3)$$

Since sum of coefficients of $P(x)$ and $Q(x)$ are equal hence $P(1) = Q(1)$. Thus $F(1) = P(1) - Q(1) = 0$.

Taking the Derivative we have

$$3(F(x))^2 F'(x) + 3(P'QF + PQ'F + PQF') = 3x^2 F'(x^3)$$

If we replace x by 1; all terms containing F will vanish. Since $P(1)$ and $Q(1)$ equals S

We have $S^2 F'(1) = F'(1)$ implying either $F'(1) = 0$ or $S^2 = 1$

If $S^2 \neq 1$ we continue the differentiation. Again ignoring all terms containing F and F' (since $F(1) = F'(1) = 0$), we have $S^2 F^2(1) = 3F^2(1)$ implying either $S^2 = 3 \vee F^2(1) = 0$.

By induction we can easily show that if we perform a times differentiation we will have

$F^a(1) = 0$ or $S^2 = 3^{a-1}$ (one 3 will be generated in each differentiation from the x^3 term; all the derivatives till $a-1$ th derivative will be ignored as at $x = 1$ they are 0.

(This is a sketch of the solution. The induction should be implemented at this step.)

Hence $F(x) = P(x) - Q(x) = (x-1)^a R(x)$ and $S^2 = 3^{a-1}$.

7. For N be a positive integer $N(N - 101)$ be a perfect square.

Since 101 is a prime number, so both N and $N - 101$ should be perfect squares.

Note that all consecutive perfect squares have a difference of successive odd numbers like

$$2^2 - 1^2 = 3$$

$$3^2 - 2^2 =$$

$$4^2 - 3^2 = 7$$

.....

$$51^2 - 50^2 = 101$$

So, $N = 51^2$ or 2601 is our required solution.

8. Since ABCD is a square so $AB \parallel CD$.

Hence the equation to the line passing through C and D must be $y = x + c$ (where c is a constant)
That is the slope is 1 (as the line AB is $y = x + 8$; it has slope 1) and the y-intercept is c .

Let the line $y = x + c$ intersect the parabola $y = x^2$ at α, β (x coordinates).

Thus $x^2 = x + c$ has the solutions α, β .

Using Sridhar's formula $x = \frac{1 \pm \sqrt{1+4c}}{2}$. These must be the values of α, β . Thus
 $\alpha - \beta = \sqrt{1+4c}$

Distance between the lines $y = x + 8$ and $y = x + c$ is $\frac{|c-8|}{\sqrt{2}}$

Also the coordinates of C and D are $(\alpha, \alpha + c), (\beta, \beta + c)$. Hence the length $CD = \sqrt{2}(\alpha - \beta)$
(using distance formula). Since $\alpha - \beta = \sqrt{1+4c}$; the length of $CD = \sqrt{2}\sqrt{1+4c}$.

This length CD equals the distance between lines $y = x + 8$ and $y = x + c$ since ABCD is a square.

Thus $\frac{|c-8|}{\sqrt{2}} = \sqrt{2}\sqrt{1+4c}$; squaring both sides we have $(c-8)^2 = 4(1+4c)$.

From here we solve for c : $c^2 - 16c + 64 = 4 + 16c$ or $c^2 - 32c + 60 = 0$
Thus $(c-30)(c-2) = 0$

Therefore c may have two values : 30 and 2

Hence the possible length of sides are $11\sqrt{2}$ or $3\sqrt{2}$
